



A formulated zinc fertilizer for foliar application.

Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

SAFETY PRECAUTIONS

STORAGE

H318: Causes serious eye damage. H410: Very toxic to aquatic life with long lasting effects. **Danger** P273: Avoid release to the environment. P280: Wear protective gloves/protective clothing/eye protection/face protection. P305+3351+338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P391: Collect spillage. P501: Dispose of contents/container as hazardous waste in accordance with local regulations. Contains: ZINC SULPHATE UFI: VE01-C39D-VN42-4H9V

Made in UK Batch No: See neck of the bottle

Contents: 10 litres

Packed in boxes of 2 x 10 litres

Manufactured by:

FMC Agro Ltd, Ŕectors Lane, Pentre, Flintshire, CH5 2DH, UK. Tel: + 44 (0) 1244 537370 - E-mail: fmc.agro.uk@fmc.com www.fmc-agro.co.uk

Imported to Ireland by: FMC Agricultural Solutions A/S Thyborønvej 78, 7673 Harboøre, Denmark. TEL:: 45 9690900

UK FERTILISER (GB only) Zinc-based fertiliser solution as sulphate Water soluble zinc (Zn): 10.7% w/w (144 g/l) Net content: 13.3 kg To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

Also contains:

Sulphur oxide (SO₃): 13% w/w

RATES OF USE AND TIMING

EC FERTILISER (Ireland only)

 Marginal deficiency:
 Apply 2.0 l/ha in at least 200 litres of water

 Moderate deficiency:
 Apply 4.0 l/ha in at least 200 litres of water

 Severe deficiency:
 Apply 4.0 l/ha in at least 200 litres of water and repeat as necessary during the growing season.

Spray whenever a zinc deficiency is observed (or expected) or as a routine maintenance treatment. Applications may commence at the 3 leaf stage or as soon as there is sufficient foliage present to absorb the spray.

(177 a/l)

Full recommendations for specific crops are available from your supplier.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

APPLICATION

Shake the container well before opening and use.

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Zinc. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

COMPATIBILITY

Zinc is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at www.fmc-agro.co.uk. Always read the label of the partner product carefully before mixing with Zinc. 11002057_22010466_GB and Ireland_12112021