



# Magnus<sup>TM</sup>

An inorganic fertiliser containing magnesium and sulphur for foliar application

## EC FERTILISER (IRELAND ONLY)

### UK FERTILISER (GB ONLY)

Suspension of magnesium hydroxide with sulphur.

Magnesium oxide (MgO): 30.0 % w/w (480 g/l)

Sulphur (S): 22.5 % w/w (360 g/l)

Net content: 16 kg

To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

## RATES OF USE AND TIMINGS

### Cereals

Apply 1 - 3 l/ha from when the crop has 3 leaves and repeat if required.

### Oilseed rape

Apply 1 - 3 l/ha in the Autumn when the crop has 6 - 8 leaves and repeat in the Spring if required.

### Peas and beans

Apply 5 l/ha when the crop is 10 - 15 cm high and repeat after 10 - 14 days if required.

### Sugar beet

Apply 2 - 4 l/ha from the 6 leaf stage. Repeat if required before the crop leaves meet across the rows.

### Potatoes

Apply 2 - 4 l/ha from when the haulm meets across the rows. Repeat if necessary.

### Maize

Apply 4 l/ha when required from the 4-leaf stage. Repeat if necessary.

### Leaf brassicas and other leafy vegetables

Apply 2 l/ha as soon as enough foliage is present to intercept the spray. Repeat if necessary.

### Root brassicas and other root vegetables

Apply 2 l/ha as soon as enough foliage is present to intercept the spray. Repeat if necessary.

## Water volumes

Use at least 200 litres of water per hectare. Increase the volume if necessary to ensure good coverage of the leaves.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

## APPLICATION

**Shake the container thoroughly before opening and use.**

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Magnus. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout. Clean sprayer thoroughly after use.

NOTE: If tank-mixing Magnus with another spray chemical, always add the partner product to the water first and make sure it is fully dispersed/dissolved before adding the Magnus to the tank.

## COMPATIBILITY

Magnus is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at [www.fmc-agro.co.uk](http://www.fmc-agro.co.uk). Always read the label of the partner product carefully before mixing with Magnus.

## STORAGE

**Protect from frost. Store above 5°C.**

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

## SAFETY PRECAUTIONS

H315: Causes skin irritation.

### Warning

P264: Wash hands thoroughly after handling.

P280: Wear protective gloves/protective clothing/eye protection/face protection.

P302+352: IF ON SKIN: Wash with plenty of water.

P332+313: If skin irritation occurs: Get medical advice/attention.

P362: Take off contaminated clothing.

Contains 1,2-Benzisothiazolin-3-one. May produce an allergic reaction.

UFI: RD21-035H-WN4F-CCQC



Made in UK

Batch No: See neck of the bottle

**Contents: 10 litres**

Manufactured by:

FMC Agro Ltd, Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.

Tel: + 44 (0) 1244 537370 - E-mail: [fmc.agro.uk@fmc.com](mailto:fmc.agro.uk@fmc.com)

[www.fmc-agro.co.uk](http://www.fmc-agro.co.uk)

Imported to Ireland by:

FMC Agricultural Solutions A/S Thybørnvej 78, 7673 Harbøre, Denmark.

TEL.: 45 96909690

FMC and Magnus are trademarks of FMC Corporation or an affiliate. ©2009-2022 FMC Corporation. All rights reserved.

