



# Thio-S™

**A high sulphur fertiliser with nitrogen for rapid correction of sulphur deficiency  
in agricultural and horticultural crops**

## Ammonium thiosulphate solution

Ammoniacal nitrogen (N): 11.6% w/w (151 g/l)

Sulphur trioxide (SO<sub>3</sub>): 64.1% w/w (862 g/l)

Net Content: 13.3 kg

To be used only where there is recognised need. Do not exceed the appropriate dose rates.

## RATES OF USE

### Cereals

Apply 3 - 5 l/ha at GS 30 - 31 and repeat at 10 - 14 day intervals if required. Wheats will benefit from a further application at GS 55 - 79.

### Oilseed rape

Apply 5 - 10 l/ha at stem extension and repeat at 10 - 14 day intervals as required.  
Do not apply during flowering.

### Potatoes

Apply 5 l/ha at or shortly after 100% emergence and repeat at 10 - 14 day intervals as required.

### Sugar beet, beans and peas

Apply 5 l/ha at the 4 - 6 leaf stage and repeat at 10 - 14 day intervals as necessary

### Market brassicas

Apply 5 l/ha when the crop has 4 - 6 leaves and repeat at 10 - 14 day intervals as required.

### Onions, leeks, carrots

Apply 5 l/ha when the crop is 15 cm tall and repeat at 10 - 14 day intervals as necessary.

### Maize

Apply 5 l/ha when the crop has 4 - 8 leaves and repeat 14 days later.

### Grass for silage

Apply 10 l/ha 4 weeks before the first (or second) cut.

### Water volume

Use in at least 200 litres of water/ha in all cases.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

## APPLICATION

Shake the container well before opening and use.

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Thio-S. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

## COMPATIBILITY

Thio-S is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at [www.fmc-agro.co.uk](http://www.fmc-agro.co.uk). Always read the label of the partner product carefully before mixing with Thio-S.

## STORAGE

**Protect from frost. Store above 5°C.**

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

## SAFETY PRECAUTIONS

The product is not classified as hazardous under GB CLP and CLP (REG EC 1272/2008).

Made in UK

Batch No: See neck of the bottle

# Contents: 10 litres

Packed in boxes of 2 x 10 litres

## Manufactured by:

FMC Agro Ltd, Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.

Tel: + 44 (0) 1244 537370 - E-mail: [fmc.agro.uk@fmc.com](mailto:fmc.agro.uk@fmc.com)

[www.fmc-agro.co.uk](http://www.fmc-agro.co.uk)

## Imported to Ireland by:

FMC Agricultural Solutions A/S Thyborønvej 78, 7673 Harboøre, Denmark.

TEL.: 45 96909690

FMC and Thio-S are trademarks of FMC Corporation or an affiliate. ©2019 FMC Corporation. All rights reserved.

